

	Pazienti (N=22)	Controlli (N=22)	Test (df)	p
<i>Protocollo PSI</i>				
HADS Tot	17,86 (±6,978)	10,77 (±5,863)	3,694 (42)	,001
HADS D	8,68 (±4,040)	4,86 (± 3,399)	3,392 (42)	,002
HADS A	8,59 (± 3,500)	5,64 (± 3,458)	2,816 (42)	,007
<i>Protocollo NPS</i>				
TMT- A	37,36 (±14,933)	33,00 (±8,635)	1,186 (33,632)	,244
TMT-B	91,05 (±38,766)	68,27 (±24,247)	2,336 (42)	,024
TMT B-A	52,77 (±30,505)	35,26 (±21,036)	2,216 (42)	,032
DIGIT F	5,00 (± 1,113)	6,05 (±0,722)	-3,697 (36,028)	,001
DIGIT B	3,95 (± 1,647)	5,05 (±1,046)	-2,623 (42)	,012
REY	48,86 (±10,246)	53,91 (± 9,232)	-1,716 (42)	,094
REY D	10,55 (8±3,291)	12,73 (± 1,142)	-2,606 (36,086)	,013
FAS	36,64 (±12,427)	47,68 (±8,091)	-3,494 (42)	,001
TOL	27 (±4,598)	31,68 (±3,872)	-3,653 (42)	,001
FACT AM	2,60 (± 0,86)	3,38 (± 0,58)	-3,507 (42)	,001
FACT CC	2,25 (± 0,86)	3,37 (± 0,84)	-4,34 0(42)	,000
FACT MEM	2,34 (±0,768)	3,23 (± 0,50)	-4,551 (36,218)	,000
FACT FV	2,20 (± 0,88)	3,33 (± 0,80)	-4,417 (42)	,000

Tabella 1 (continua nella pag. successiva). Risultati del confronto fra medie di gruppo al t-test

	Pazienti (N=22)	Controlli (N=22)	Test (df)	p
FACT ATT	2,33 (± 0,71)	3,17 (±0,55)	-4,310 (42)	,000
FACT OE	2,47(±0,97)	3,37 (± 0,49)	-3,853 (31,015)	,001
FACT QoL	2,09 (± 1,07)	3,5 (± 0,40)	-5,953 (26,730)	,001
FACT TOT	2,15 (± 0,63)	2,97 (± 0,36)	-5,218 (33,368)	,000
<i>Protocollo Emozioni</i>				
EKMAN Tot	47,73 (±5,035)	53,05 (±3,031)	-4,244 (34,455)	,000
EKMAN Rabbia	7,59 (±1,436)	8,55 (± 1,262)	-2,342 (42)	,024
EKMAN Tristezza	7,45 (±1,738)	7,86 (±1,167)	-,917 (36,731)	,365
EKMAN Sorpresa	8,92 (±1,065)	9,45 (±0,800)	-1,920 (42)	,062
EKMAN Paura	6,00 (± 2,600)	8,09 (± 1,411)	-3,315 (32,382)	,002
EKMAN Disgusto	8,05(± 1,582)	9,50 (± 0,802)	-3,835 (42)	,000
EKMAN Felicità	9,77 (±0,528)	9,59 (±0,796)	,892 (42)	,377
RME Controllo	35,00 (±1,155)	35,32 (± 0,716)	-1,098 (42)	,278
RME Sperimentale	24,86 (±4,201)	27,91 (±2617)	-2,886 (42)	,006
EQ	45,09 (±9,092)	52,32 ± (8,277)	-2,757 (42)	,009
TAS Tot	51,41 (±8,700)	41,14 (±8,806)	3,892 (42)	,000
TAS F1	19,32 (±6,643)	14,55 (±4667)	2,757 (42)	,009
TAS F2	14,45 (±4,394)	11,09 (±3,939)	2,674 (42)	,011
TAS F3	17,68 (±3,695)	15,50 (±4,798)	1,690 (42)	,099

Tabella 1. Risultati del confronto fra medie di gruppo al t-test

	Pazienti (N=22) Media (ds)	Controlli (N=22) Media (ds)	p Scolarità	p Pazienti
<i>Protocollo NPS</i>				
TMT- A	37,36 (±14,933)	33,00 (±8,635)	,002	,242
TMT-B	91,05 (±38,766)	68,27 (±24,247)	,038	,657
TMT B-A	52,77 (±30,505)	35,26 (±21,036)	,232	,374
DIGIT F	5,00 (± 1,113)	6,05 (±0,722)	,921	,007
DIGIT B	3,95 (± 1,647)	5,05 (±1,046)	,531	,022
REY	48,86 (±10,246)	53,91 (± 9,232)	,135	,746
REY D	10,55 (8±3,291)	12,73 (± 1,142)	,092	,376
FAS	36,64 (±12,427)	47,68 (±8,091)	,033	,192
TOL	27 (±4,598)	31,68 (±3,872)	,494	,027
<i>Protocollo Emozioni</i>				
EKMAN Tot	47,73 (±5,035)	53,05 (±3,031)	,117	,031
RME Sperimentale	24,86 (±4,201)	27,91 (±2617)	,029	,419
EQ	45,09 (±9,092)	52,32 ± (8,277)	,640	,085

Tabella 2. significatività della differenza fra medie nei due gruppi per le variabili potenzialmente influenzabili dal livello di scolarità